



Contribution ID: 65

Type: **Contributed talk**

## Galactic chemical Evolution with short lived radioactive isotopes

*Tuesday 17 September 2024 15:05 (15 minutes)*

Studying the galactic chemical evolution with short lived radioisotopes (SLRs) has a significant advantage over using stable elements: Due to their radioactive decay, SLRs carry additional timing information on astrophysical nucleosynthesis sites.

We can use meteoritic abundance data in conjunction with a chemical evolution model to constrain the physical conditions in the last rapid neutron capture process event that polluted the early Solar system prior to its formation [1].

Further, with the help of detections of live SLRs of cosmic origin in the deep sea crust [2], we can use these data in a 3-dimensional chemical evolution code to explain why different classes of radioisotopes should often arrive conjointly on Earth, even if they were produced in different sites (e.g., neutron star mergers, core-collapse/thermonuclear supernovae) [3].

Finally, we included radioisotope production into a cosmological zoom-in simulation to create a map of Al-26 indicating areas of ongoing star formation in the Galaxy, consistent with the observations by the SPI/INTEGRAL instrument[4]. We provide predictions for future gamma-ray detection instruments.

### References:

- [1] Côté et al., 2021 Science 371, 945
- [2] Wallner et al., 2021 Science 372, 742W
- [3] Wehmeyer et al., 2023 ApJ 944, 121
- [4] Kretschmer et al., 2013 A&A 559, A9

**Primary author:** WEHMEYER, Benjamin (University of Wroclaw)

**Co-authors:** Dr YAGÜE LÓPEZ, Andrés (Los Alamos National Laboratory); Dr CÔTÉ, Benoit (University of Victoria); Dr K. PETŐ, Maria (Konkoly Observatory); Prof. KOBAYASHI, Chiaki (University of Hertfordshire); LUGARO, Maria (Konkoly Observatory, CSFK HUN-REN)

**Presenter:** WEHMEYER, Benjamin (University of Wroclaw)

**Session Classification:** Plenary Session